

ANNEX 2B

MANUFACTURE OF BIOLOGICAL MEDICINAL SUBSTANCES AND PRODUCTS FOR HUMAN USE

SCOPE

The methods employed in the manufacture of biological active substances and biological medicinal products for human use ('biological active substances and medicinal products') are a critical factor in shaping the appropriate regulatory control. Biological active substances and medicinal products can be defined therefore largely by reference to their method of manufacture. This annex provides guidance on the full range of active substances and medicinal products defined as biological with the exception of Advanced Therapy Medicinal Products ("ATMPs"). The ATMPs are not covered by the present guideline. Manufacturers of ATMPs should refer to PIC/S Annex 2A Manufacture of Advanced Therapy Medicinal Products for Human Use.

This annex is divided into two main parts:

- a) Part A contains supplementary guidance on the manufacture of biological active substances and medicinal products, from control over seed lots and cell banks through to finishing activities and testing.
- b) Part B contains further guidance on selected types of biological active substances and medicinal products.

This annex, along with several other annexes of the PIC/S Guide to GMP, provides guidance which supplements that in Part I and in Part II of the Guide. There are two aspects to the scope of this annex:

- a) Stage of manufacture - for biological active substances to the point immediately prior to their being rendered sterile, the primary guidance source is Part II. Guidance for the subsequent manufacturing steps of biological products are covered in Part I.
- b) Type of product - this annex provides guidance on the full range of medicinal products defined as biological with the exception of ATMPs.

These two aspects are shown in Table 1; it should be noted that this table is illustrative only and is not meant to describe the precise scope. It should also be understood that in line with the corresponding table in Part II of the Guide, the level of GMP increases in detail from early to later steps in the manufacture of biological active substances but GMP principles should always be adhered to. The inclusion of some early steps of manufacture within the scope of this Annex does not imply that those steps will be routinely subject to inspection by the authorities.

47 Antibiotics are not defined as biological medicinal products, however where
 48 biological stages of manufacture occur, guidance in this Annex may be used.

49
 50 Guidance for medicinal products derived from fractionated human blood or
 51 plasma is covered in Annex 14 and for non-transgenic plant products in Annex 7.

52
 53 In certain cases, other legislation may be applicable to the starting materials for
 54 biologicals. For example,

55 (a) Tissue and cells used as starting materials for medicinal products,
 56 donation, procurement, testing, processing, preservation, storage and
 57 distribution of human tissues and cells of tissue and cells may be covered
 58 by national legislation. Such tissues and cells may provide the active
 59 substances for some biological medicinal product within the scope of this
 60 annex at which point GMP and other medicinal product legislation
 61 requirements apply.

62 (b) Blood or blood components used as starting materials for medicinal
 63 products, national legislation may provide the technical requirements for
 64 the selection of donors, collection, testing, processing, storage, and
 65 distribution of human blood and blood components¹.

66 Additionally, the manufacture and control of genetically modified organisms
 67 needs to comply with local and national requirements. Appropriate containment
 68 should be established and maintained in facilities where any genetically modified
 69 micro-organism is handled². Advice should be obtained according to national
 70 legislation in order to establish and maintain the appropriate Biological Safety
 71 Level. There should be no conflicts with GMP requirements.

72
 73
 74

Table 1. Illustrative guide to manufacturing activities within the scope of Annex 2B

Type and source of material	Example product	Application of this guide to manufacturing steps shown in grey			
1. Animal or plant sources: non-transgenic	Heparins, insulin, enzymes, proteins, allergen extract, immunosera	Collection of plant, organ, animal material or fluid ³	Cutting, mixing, and / or initial processing	Isolation and purification	Formulation, filling
2. Virus or bacteria / fermentation / cell culture	Viral or bacterial vaccines; enzymes, proteins	Establishment & maintenance of MCB ⁴ , WCB, MVS, WVS	Cell culture and/or fermentation	Inactivation when applicable, isolation and purification	Formulation, filling
3. Biotechnology fermentation/ cell culture	Recombinant products, MAb, allergens, vaccines	Establishment & maintenance of MCB and WCB, MSL, WSL	Cell culture and /or fermentation	Isolation, purification, modification	Formulation, filling


¹ In the EEA, this is Directive 2002/98/EC and its Commission Directives.

² In the EEA, this is Directive 2009/41/EC on contained use of genetically modified micro-organisms.

³ See section B1 for the extent to which GMP principles apply.

⁴ See section on 'Seed lot and cell bank system' for the extent to which GMP applies.

4. Animal sources: transgenic	Recombinant proteins	Master and working transgenic bank	Collection, cutting, mixing, and/or initial processing	Isolation, purification and modification	Formulation, filling
5. Plant sources: transgenic	Recombinant proteins, vaccines, allergens	Master and working transgenic bank	Growing, harvesting ⁵	Initial extraction, isolation, purification, modification	Formulation, filling
6. Human sources	Urine derived enzymes, hormones	Collection of fluid ⁶	Mixing, and/or initial processing	Isolation and purification	Formulation, filling
7. Human sources	Products from cells and tissues	Donation, procurement and testing of starting tissue / cells ⁷	Initial processing, isolation and purification.	Cell isolation, culture, purification, combination with non-cellular components	Formulation, combination, filling

75
76
77

78

79 See Glossary for explanation of acronyms.

80

81 **PRINCIPLE**

82

83 The manufacture of biological active substances and medicinal products involves
84 certain specific considerations arising from the nature of the products and the
85 processes. The ways in which biological medicinal products are manufactured,
86 controlled and administered make some particular precautions necessary.

87

88 Unlike conventional medicinal products, which are manufactured using chemical
89 and physical techniques capable of a high degree of consistency, the
90 manufacture of biological active substances and medicinal products involves
91 biological processes and materials, such as cultivation of cells or extraction from
92 living organisms. These biological processes may display inherent variability, so
93 that the range and nature of by-products may be variable. As a result, quality risk
94 management (QRM) principles are particularly important for this class of
95 materials and should be used to develop the control strategy across all stages of
96 manufacture so as to minimise variability and to reduce the opportunity for
97 contamination and cross-contamination.

98

99 Since materials and processing conditions used in cultivation processes are
100 designed to provide conditions for the growth of specific cells and
101 microorganisms, this provides extraneous microbial contaminants the opportunity

⁵ In the EEA: HMPC guideline on Good Agricultural and Collection Practice - EMEA/HMPC/246816/2005 may be applied to growing, harvesting and initial processing in open fields.

⁶ For principles of GMP apply, see explanatory text in 'Scope'.

⁷ In the EEA, human tissues and cells must comply with Directive 2004/23/EC and implementing Directives at these stages.

102 to grow. In addition, some products may be limited in their ability to withstand a
103 wide range of purification techniques particularly those designed to inactivate or
104 remove adventitious viral contaminants. The design of the processes, equipment,
105 facilities, utilities, the conditions of preparation and addition of buffers and
106 reagents, sampling and training of the operators are key considerations to
107 minimise such contamination events.

108
109 Specifications related to products (such as those in Pharmacopoeial
110 monographs, Clinical Trial Authorisation (CTA), and Marketing Authorisation
111 (MA)) will dictate whether and to what stage substances and materials can have
112 a defined level of bioburden or need to be sterile. Similarly, manufacturing must
113 be consistent with other specifications set out in the CTA or MA (e.g. number of
114 generations (doublings, passages) between the seed lot or cell bank).

115
116 For biological materials that cannot be sterilized (e.g. by filtration), processing
117 must be conducted aseptically to minimise the introduction of contaminants.
118 Where they exist, other guidance documents should be consulted on the
119 validation of specific manufacturing methods, e.g. virus removal or inactivation.
120 The application of appropriate environmental controls and monitoring and,
121 wherever feasible, in-situ cleaning and sterilisation systems together with the use
122 of closed systems can significantly reduce the risk of accidental contamination
123 and cross-contamination.

124
125 Control usually involves biological analytical techniques, which typically have a
126 greater variability than physico-chemical determinations. A robust manufacturing
127 process is therefore crucial and in-process controls take on a particular
128 importance in the manufacture of biological active substances and medicinal
129 products.

130
131 Biological medicinal products which incorporate human tissues or cells must
132 comply with national requirements for the coding, processing, preservation,
133 storage and distribution of human tissues and cells.⁸ Collection and testing of this
134 material must be done in accordance with an appropriate quality system and in
135 accordance with applicable national requirements⁹. Furthermore, national
136 requirements¹⁰ on traceability apply from the donor (while maintaining donor
137 confidentiality) through stages applicable at the Tissue Establishment and then
138 continued under medicines legislation through to the institution where the product
139 is used.

140
141 Biological active substances and medicinal products must comply with the
142 applicable national guidance on minimising the risk of transmitting animal
143 spongiform encephalopathy agents via human and veterinary medicinal
144 products.

145
146

⁸ In the EEA, these are Directive 2004/23/EC and Directive 2006/17/EC.

⁹ In the EEA, this is the Commission Directive 2006/86/EC.

¹⁰ In the EEA, this is Directive 2006/86/EC.

PART A. GENERAL GUIDANCE

PERSONNEL

1. Personnel (including those concerned with cleaning, maintenance or quality control) employed in areas where biological active substances and products are manufactured and tested should receive training, and periodic retraining, specific to the products manufactured and to their work, including any specific security measures to protect product, personnel and the environment.
2. The health status of personnel should be taken into consideration for product safety. Where necessary, personnel engaged in production, maintenance, testing and animal care (and inspections) should be vaccinated with appropriate specific vaccines and have regular health checks.
3. Any changes in the health status of personnel, which could adversely affect the quality of the product, should preclude work in the production area and appropriate records kept. Production of BCG vaccine and tuberculin products should be restricted to staff who are carefully monitored by regular checks of immunological status or chest X-ray. Health monitoring of staff should be commensurate with the risk, medical advice should be sought for personnel involved with hazardous organisms.
4. Where required to minimise the opportunity for cross-contamination, restrictions on the movement of all personnel (including quality control (QC), maintenance and cleaning staff) should be controlled on the basis of QRM principles. In general, personnel should not pass from areas where exposure to live micro-organisms, genetically modified organisms, toxins or animals to areas where other products, inactivated products or different organisms are handled. If such passage is unavoidable, the contamination control measures should be based on QRM principles.

PREMISE AND EQUIPMENT

5. As part of the control strategy, the degree of environmental control of particulate and microbial contamination of the production premises should be adapted to the active substance, intermediate or finished product and the production step, bearing in mind the potential level of contamination of the starting materials and the risks to the product. The environmental monitoring programme should be supplemented by the inclusion of methods to detect the presence of specific microorganisms (i.e. host organism, yeasts, moulds, anaerobes, etc) where indicated by the QRM process.
6. Manufacturing and storage facilities, processes and environmental classifications should be designed to prevent the extraneous contamination of products. Prevention of contamination is more appropriate than detection and removal, although contamination is likely to become evident during processes such as fermentation and cell culture. Where processes are not closed and there is therefore exposure of the product to the immediate room environment (e.g. during additions of supplements, media, buffers, gasses,) control measures should be put in place, including engineering and environmental controls on the basis of

- 199 QRM principles. These QRM principles should take into account the principles
200 and guidance from the appropriate sections of Annex 1¹¹ when selecting
201 environmental classification cascades and associated controls.
202
- 203 7. Dedicated production areas should be used for the handling of live cells, capable
204 of persistence in the manufacturing environment,. Dedicated production area
205 should be used for the manufacture of pathogenic organisms (i.e. Biosafety level
206 3 or 4).
207
- 208 8. Manufacture in a multi-product facility may be acceptable where the following, or
209 equivalent (as appropriate to the product types involved) considerations and
210 measures are part of an effective control strategy to prevent cross-contamination:
- 211 (a) Knowledge of key characteristics of all cells, organisms and any
212 adventitious agents (e.g. pathogenicity, detectability, persistence,
213 susceptibility to inactivation) within the same facility.
- 214 (b) Where production is characterised by multiple small batches from different
215 starting materials, factors such as the health status of donors and the risk
216 of total loss of product should be taken into account when considering the
217 acceptance of concurrent working during development of the control
218 strategy.
- 219 (c) Live organisms and spores are prevented from entering non-related areas
220 or equipment by addressing all potential routes of cross-contamination and
221 utilizing single use components and engineering measures such as closed
222 systems.
- 223 (d) Control measures to remove the organisms and spores before the
224 subsequent manufacture of other products, these control measures should
225 also take the heating, ventilation and air conditioning (HVAC) system into
226 account. Cleaning and decontamination for the organisms and spores
227 should be validated.
- 228 (e) Environmental monitoring, specific for the micro-organism being
229 manufactured, where the micro-organisms are capable of persistence in
230 the manufacturing environment and where methods are available, is
231 conducted in adjacent areas during manufacture and after completion of
232 cleaning and decontamination. Attention should also be given to risks
233 arising with use of certain monitoring equipment (e.g. airborne particle
234 monitoring) in areas handling live and/or spore forming organisms.
- 235 (f) Products, equipment, ancillary equipment (e.g. for calibration and
236 validation) and disposable items are only moved within and removed from
237 such areas in a manner that prevents contamination of other areas, other
238 products and different product stages (e.g. prevent contamination of
239 inactivated or toxoided products with non-inactivated products).
- 240 (g) Campaign based manufacturing.
241

¹¹ Although the title of Annex 1 refers to the manufacture of sterile medicinal products it is not the intention to force the manufacture of sterile product at a stage when a low bioburden is appropriate and authorised. Its use is because it is the PIC/S GMP source of guidance on all of the classified manufacturing areas including the lower grades D and C.

- 242 9. For finishing (secondary) operations¹², the need for dedicated facilities will
243 depend on consideration of the above together with additional considerations
244 such as the specific needs of the biological medicinal product and on the
245 characteristics of other products, including any non-biological products, in the
246 same facility. Other control measures for finishing operations may include the
247 need for specific addition sequences, mixing speeds, time and temperature
248 controls, limits on exposure to light and containment and cleaning procedures in
249 the event of spillages.
250
- 251 10. The measures and procedures necessary for containment (i.e. for environment
252 and operator safety) should not conflict with those for product quality.
253
- 254 11. Air handling units should be designed, constructed and maintained to minimise
255 the risk of cross-contamination between different manufacturing areas and may
256 need to be specific for an area. Consideration, based on QRM principles, should
257 be given to the use of single pass air systems.
258
- 259 12. Positive pressure areas should be used to process sterile products but negative
260 pressure in specific areas at the point of exposure of pathogens is acceptable for
261 containment reasons. Where negative pressure areas or safety cabinets are used
262 for aseptic processing of materials with particular risks (e.g. pathogens), they
263 should be surrounded by a positive pressure clean zone of appropriate grade.
264 These pressure cascades should be clearly defined and continuously monitored
265 with appropriate alarm settings.
266
- 267 13. Equipment used during handling of live organisms and cells, including those for
268 sampling, should be designed to prevent any contamination during processing.
269
- 270 14. Primary containment¹³ should be designed and periodically tested to ensure the
271 prevention of escape of biological agents into the immediate working
272 environment.
273
- 274 15. The use of 'clean in place' and 'steam in place' ('sterilisation in place') systems
275 should be used where possible. Valves on fermentation vessels should be
276 completely steam sterilisable.
277
- 278 16. Air vent filters should be hydrophobic and validated for their scheduled life span
279 with integrity testing at appropriate intervals based on appropriate QRM
280 principles.
281
- 282 17. Drainage systems must be designed so that effluents can be effectively
283 neutralised or decontaminated to minimise the risk of cross-contamination. Local
284 regulation must be complied with to minimise the risk of contamination of the
285 external environment according to the risk associated with the biohazardous
286 nature of waste materials.
287
- 288 18. Due to the variability of biological products or manufacturing processes,
289 relevant/critical raw materials (such as culture media and buffers) have to be
290 measured or weighed during the production process. In these cases, small stocks

¹² Formulation, filling and packaging

¹³ See main GMP Glossary on 'Containment'.

291 of these raw materials may be kept in the production area for a specified duration
292 based on defined criteria such as for the duration of manufacture of the batch or
293 of the campaign.
294

295

296 ANIMALS

297

298 19. A wide range of animal species are used in the manufacture of a number of
299 biological medicinal products. These can be divided into 2 broad types of
300 sources:

301 (a) Live groups, herds, flocks: examples include polio vaccine (monkeys),
302 immunosera to snake venoms and tetanus (horses, sheep and goats),
303 allergens (cats), rabies vaccine (rabbits, mice and hamsters), transgenic
304 products (goats, cattle).

305 (b) Animal materials derived post-mortem and from establishments such as
306 abattoirs: examples include, abattoir sources for enzymes, anticoagulants
307 and hormones (sheep and pigs).

308 In addition, animals may also be used in quality control either in generic assays,
309 e.g. pyrogenicity, or specific potency assays, e.g. pertussis vaccine (mice),
310 pyrogenicity (rabbits), BCG vaccine (guinea-pigs).
311

311

312 20. In addition to compliance with TSE regulations, other adventitious agents that are
313 of concern (zoonotic diseases, diseases of source animals) should be monitored
314 by an ongoing health programme and recorded. Specialist advice should be
315 obtained in establishing such programmes. Instances of ill-health occurring in the
316 source/donor animals should be investigated with respect to their suitability and
317 the suitability of in-contact animals for continued use (in manufacture, as sources
318 of starting and raw materials, in quality control and safety testing), the decisions
319 must be documented. A look-back procedure should be in place which informs
320 the decision making process on the continued suitability of the biological active
321 substance or medicinal product in which the animal sourced starting or raw
322 materials have been used or incorporated. This decision-making process may
323 include the re-testing of retained samples from previous collections from the
324 same donor animal (where applicable) to establish the last negative donation.
325 The withdrawal period of therapeutic agents used to treat source/donor animals
326 must be documented and used to determine the removal of those animals from
327 the programme for defined periods.
328

328

329 21. Particular care should be taken to prevent and monitor infections in the source /
330 donor animals. Measures should include the sourcing, facilities, husbandry,
331 biosecurity procedures, testing regimes, control of bedding and feed materials.
332 This is of special relevance to specified pathogen free animals where
333 pharmacopoeial monograph requirements must be met. Housing and health
334 monitoring should be defined for other categories of animals (e.g. healthy flocks
335 or herds).
336

336

337 22. For products manufactured from transgenic animals, traceability should be
338 maintained in the creation of such animals from the source animals.
339

339

- 340 23. Note should be taken of national requirements on the protection of animals used
341 for scientific purposes¹⁴. Housing for animals used in production and control of
342 biological active substances and medicinal products should be separated from
343 production and control areas.
344
- 345 24. For different animal species, key criteria should be defined, monitored, and
346 recorded. These may include age, weight and health status of the animals.
347
- 348 25. Animals, biological agents, and tests carried out should be the subject of an
349 identification system to prevent any risk of confusion and to control all identified
350 hazards.
351
352

353 DOCUMENTATION

- 354
- 355 26. Starting and raw materials may need additional documentation on the source,
356 origin, distribution chain, method of manufacture, and controls applied, to assure
357 an appropriate level of control including their microbiological quality.
358
- 359 27. Some product types may require specific definition of what materials constitutes
360 a batch, particularly cells. For autologous and donor-matched situations, the
361 manufactured product should be viewed as a batch.
362
- 363 28. Where human cell or tissue donors are used, full traceability is required from
364 starting and raw materials, including all substances coming into contact with the
365 cells or tissues through to confirmation of the receipt of the products at the point
366 of use whilst maintaining the privacy of individuals and confidentiality of health
367 related information¹⁵. Traceability records must be retained for 30 years after the
368 expiry date of the medicinal product. Particular care should be taken to maintain
369 the traceability of products for special use cases, such as donor-matched cells.
370 National requirements¹⁶ in regards to traceability requirements and notification of
371 serious adverse reactions and events apply to blood components when they are
372 used as starting or raw materials in the manufacturing process of medicinal
373 products.
374
375

376 PRODUCTION

- 377
- 378 29. Given the variability inherent in many biological active substances and medicinal
379 products, steps to increase process robustness thereby reducing process
380 variability and enhancing reproducibility at the different stages of the product
381 lifecycle such as process design should be reassessed during Product Quality
382 Reviews.
383
- 384 30. Since cultivation conditions, media and reagents are designed to promote the
385 growth of cells or microbial organisms, typically in an axenic state, particular
386 attention should be paid in the control strategy to ensure there are robust steps
387 that prevent or minimise the occurrence of unwanted bioburden and associated

¹⁴ In the EEA, this is Directive 2010/63/EC.

¹⁵ In the EEA, see Article 15 of Regulation 1394/ 2007.

¹⁶ In the EEA, these are Directives 2002/98/EC and 2005/61/EC.

388 metabolites and endotoxins. For medicinal products from cells and tissues where
389 production batches are frequently small the risk of cross-contamination between
390 cell preparations from different donors with various health status should be
391 controlled under defined procedures and requirements.

392
393

394 **STARTING AND RAW MATERIALS**

395

396 31. The source, origin and suitability of biological starting and raw materials (e.g.
397 cryoprotectants, feeder cells, reagents, culture media, buffers, serum, enzymes,
398 cytokines, growth factors) should be clearly defined. Where the necessary tests
399 take a long time, it may be permissible to process starting materials before the
400 results of the tests are available, the risk of using a potentially failed material and
401 its potential impact on other batches should be clearly understood and assessed
402 under the principles of QRM. In such cases, release of a finished product is
403 conditional on satisfactory results of these tests. The identification of all starting
404 materials should be in compliance with the requirements appropriate to its stage
405 of manufacture. For biological medicinal products further guidance can be found
406 in Part I and Annex 8 and for biological active substances in Part II.

407

408 32. The risk of contamination of starting and raw materials during their passage along
409 the supply chain must be assessed, with particular emphasis on TSE. Materials
410 that come into direct contact with manufacturing equipment or the product (such
411 as media used in media fill experiments and lubricants that may contact the
412 product) must also be taken into account.

413

414 33. Given that the risks from the introduction of contamination and the consequences
415 to the finished product is the same irrespective of the stage of manufacture,
416 establishment of a control strategy to protect the product and the preparation of
417 solutions, buffers and other additions should be based on the principles and
418 guidance contained in the appropriate sections of Annex 1. The controls required
419 for the quality of starting and raw materials and on the aseptic manufacturing
420 process, assume greater importance particularly for products, in respect of which
421 final sterilisation is not possible. Where a CTA or MA provides for an allowable
422 type and level of bioburden, for example at active substance stage, the control
423 strategy should address the means by which this is maintained within the
424 specified limits.

425

426 34. Where sterilisation of starting and raw materials is required, it should be carried
427 out where possible by heat. Where necessary, other appropriate methods may
428 also be used for inactivation of biological materials (e.g. irradiation and filtration).

429

430 35. Reduction in bioburden associated with procurement of living tissues and cells
431 may require the use of other measures such as antibiotics at early manufacturing
432 stages. This should be avoided, but where it is necessary their use should be
433 justified, they should be removed from the manufacturing process at the stage
434 specified in the CTA or MA.

435

- 436 36. The donation, procurement and testing of human tissues and cells used as
437 starting materials for biological medicinal products should be in accordance with
438 national law.¹⁷ Traceability for human tissues and cells used as starting materials
439 for biological medicinal products should be maintained from the donor to the
440 batch of a finished medicinal product. Appropriate arrangements should be made
441 between the manufacturer and the supplier of tissues and cells regarding the
442 transfer of health donor information that may become available after the supply
443 of the starting material and which may have an impact on the quality or safety of
444 the medicinal product manufactured therefrom.
- 445 (a) Their procurement, donation and testing is regulated in some countries¹⁸.
446 Such supply sites must hold appropriate approvals from the national
447 competent authority(ies) which should be verified as part of starting material
448 supplier management.
- 449 (b) Where such human cells or tissues are imported they must meet equivalent
450 national standards of quality and safety¹⁹. The traceability and serious
451 adverse reaction and serious adverse event notification requirements may
452 be set out in national legislation²⁰.
- 453 (c) There may be some instances where processing of cells and tissues used
454 as starting materials for biological medicinal products will be conducted at
455 tissue establishments²¹.
- 456 (d) Tissue and cells are released by the Responsible Person (RP) in the tissue
457 establishment before shipment to the medicinal product manufacturer, after
458 which normal medicinal product starting material controls apply. The test
459 results of all tissues / cells supplied by the tissue establishment should be
460 available to the manufacturer of the medicinal product. Such information
461 must be used to make appropriate material segregation and storage
462 decisions. In cases where manufacturing must be initiated prior to receiving
463 test results from the tissue establishment, tissue and cells may be shipped
464 to the medicinal product manufacturer provided controls are in place to
465 prevent cross-contamination with tissue and cells that have been released
466 by the RP in the tissue establishment.
- 467 (e) The transport of human tissues and cells to the manufacturing site must be
468 controlled by a written agreement between the responsible parties. The
469 manufacturing sites should have documentary evidence of adherence to
470 the specified storage and transport conditions.
- 471 (f) Continuation of traceability requirements started at tissue establishments
472 through to the recipient(s), and vice versa, including materials in contact
473 with the cells or tissues, should be maintained.
- 474 (g) A technical agreement should be in place between the responsible parties
475 (e.g. manufacturers, tissue establishment, Sponsors, MA Holder) which
476 defines the tasks of each party, including the RP and Authorised Person.

¹⁷ In the EEA, this is Directive 2004/23/EC or for blood-derived cells, compliance with Directive 2002/98 regarding donation, procurement and testing.

¹⁸ In the EEA, this is Directive 2004/23/EC and its Commission directives.

¹⁹ In the EEA, they must be equivalent to those laid down in Directive 2004/23/EC.

²⁰ In the EEA, this is Directive 2006/86/EC.

²¹ In the EEA, such processing steps, are under the scope of 2004/23/EC and the Responsible Person (RP).

- 477
478 37. (...) ²²
479
480 38. Where human or animal cells are used in the manufacturing process as feeder
481 cells, appropriate controls over the sourcing, testing, transport and storage
482 should be in place ²³, including control of compliance with national requirements
483 for human cells.
484
485

486 SEED LOT AND CELL BANK SYSTEM

487
488

- 489 39. In order to prevent the unwanted drift of properties which might ensue from
490 repeated subcultures or multiple generations, the production of biological
491 medicinal substances and products obtained by microbial culture, cell culture or
492 propagation in embryos and animals should be based on a system of master and
493 working virus seed lots and/or cell banks.
494
495 40. The number of generations (doublings, passages) between the seed lot or cell
496 bank, the biological active substance and the finished product should be
497 consistent with specifications in the CTA or MA.
498
499 41. As part of product lifecycle management, establishment of seed lots and cell
500 banks, including master and working generations, should be performed under
501 circumstances which are demonstrably appropriate. This should include an
502 appropriately controlled environment to protect the seed lot and the cell bank and
503 the personnel handling it. During the establishment of the seed lot and cell bank,
504 no other living or infectious material (e.g. virus, cell lines or cell strains) should
505 be handled simultaneously in the same area or by the same persons. For stages
506 prior to the master seed or cell bank generation, where only the principles of GMP
507 may be applied, documentation should be available to support traceability
508 including issues related to components used during development with potential
509 impact on product safety (e.g. reagents of biological origin) from initial sourcing
510 and genetic development if applicable. For vaccines the requirements of
511 pharmacopoeial monographs will apply ²⁴.
512
513 42. Following the establishment of master and working cell banks and master and
514 working seed lots, quarantine and release procedures should be followed. This
515 should include adequate characterization and testing for contaminants. Their
516 on-going suitability for use should be further demonstrated by the consistency of
517 the characteristics and quality of the successive batches of product. Evidence of
518 the stability and recovery of the seeds and banks should be documented and
519 records should be kept in a manner permitting trend evaluation.
520
521 43. Seed lots and cell banks should be stored and used in such a way as to minimize
522 the risks of contamination (e.g. stored in the vapour phase of liquid nitrogen in
523 sealed containers) or alteration. Ensuring compliance with measures for the

²² This line has been intentionally left blank to harmonise with the formatting structure of the EU GMP Guide.

²³ In the EEA, this includes compliance with Directive 2004/23 EC for human cells.

²⁴ In the EEA, this is Ph Eur monograph 2005;153 "Vaccines for human use".

- 524 storage of different seeds and/or cells in the same area or equipment should
525 prevent mix-up and take into account the infectious nature of the materials to
526 prevent cross contamination.
527
- 528 44. This line has been intentionally left blank to harmonize with the structure of a
529 PIC/S Partner.
530
- 531 45. Storage containers should be sealed, clearly labelled and kept at an appropriate
532 temperature. A stock inventory must be kept. The storage temperature should be
533 recorded continuously and, where used, the liquid nitrogen level monitored.
534 Deviation from set limits and corrective and preventive action taken should be
535 recorded.
536
- 537 46. It is desirable to split stocks and to store the split stocks at different locations so
538 as to minimize the risks of total loss. The controls at such locations should provide
539 the assurances outlined in the preceding paragraphs.
540
- 541 47. The storage and handling conditions for stocks should be managed according to
542 the same procedures and parameters. Once containers are removed from the
543 seed lot / cell bank management system, the containers should not be returned
544 to stock.
545
546

547 **OPERATING PRINCIPLES**

- 548
- 549 48. Change management should, on a periodic basis, take into account the effects,
550 including cumulative effects of changes (e.g. to the process) on the quality, safety
551 and efficacy of the finished product.
552
- 553 49. Critical operational (process) parameters, or other input parameters which affect
554 product quality, need to be identified, validated, documented and be shown to be
555 maintained within requirements.
556
- 557 50. A control strategy for the entry of articles and materials into production areas
558 should be based on QRM principles. For aseptic processes, heat stable articles
559 and materials entering a clean area or clean/contained area should preferably do
560 so through a double-ended autoclave or oven. Heat labile articles and materials
561 should enter through an air lock with interlocked doors where they are subject to
562 effective surface sanitisation procedures. Sterilisation of articles and materials
563 elsewhere is acceptable provided that they are multiple wrappings, as
564 appropriate to the number of stages of entry to the clean area, and enter through
565 an airlock with the appropriate surface sanitisation precautions.
566
- 567 51. The growth promoting properties of culture media should be demonstrated to be
568 suitable for its intended use. If possible, media should be sterilized in situ. In-line
569 sterilizing filters for routine addition of gases, media, acids or alkalis, anti-foaming
570 agents etc. to fermenters should be used where possible.
571
- 572 52. Addition of materials or cultures to fermenters and other vessels and sampling
573 should be carried out under carefully controlled conditions to prevent
574 contamination. Care should be taken to ensure that vessels are correctly
575 connected when addition or sampling takes place.

- 576
577 53. Continuous monitoring of some production processes (e.g. fermentation) may be
578 necessary; such data should form part of the batch record. Where continuous
579 culture is used, special consideration should be given to the quality control
580 requirements arising from this type of production method.
581
- 582 54. Centrifugation and blending of products can lead to aerosol formation and
583 containment of such activities to minimise cross-contamination is necessary.
584
- 585 55. Accidental spillages, especially of live organisms, must be dealt with quickly and
586 safely. Qualified decontamination measures should be available for each
587 organism or groups of related organisms. Where different strains of single
588 bacteria species or very similar viruses are involved, the decontamination
589 process may be validated with one representative strain, unless there is reason
590 to believe that they may vary significantly in their resistance to the agent(s)
591 involved.
592
- 593 56. If obviously contaminated, such as by spills or aerosols, or if a potential
594 hazardous organism is involved, production and control materials, including
595 paperwork, must be adequately disinfected, or the information transferred out by
596 other means.
597
- 598 57. In cases where a virus inactivation or removal process is performed during
599 manufacture, measures should be taken to avoid the risk of recontamination of
600 treated products by non-treated products.
601
- 602 58. For products that are inactivated by the addition of a reagent (e.g.
603 micro-organisms in the course of vaccine manufacture) the process should
604 ensure the complete inactivation of live organism. In addition to the thorough
605 mixing of culture and inactivant, consideration should be given to contact of all
606 product-contact surfaces exposed to live culture and, where required, the transfer
607 to a second vessel.
608
- 609 59. A wide variety of equipment is used for chromatography. QRM principles should
610 be used to devise the control strategy on matrices, the housings and associated
611 equipment when used in campaign manufacture and in multi-product
612 environments. The re-use of the same matrix at different stages of processing is
613 discouraged. Acceptance criteria, operating conditions, regeneration methods,
614 life span and sanitization or sterilisation methods of columns should be defined.
615
- 616 60. Where irradiated equipment and materials are used, Annex 12 should be
617 consulted for further guidance.
618
- 619 61. There should be a system to assure the integrity and closure of containers after
620 filling where the final products or intermediates represent a special risk and
621 procedures to deal with any leaks or spillages. Filling and packaging operations
622 need to have procedures in place to maintain the product within any specified
623 limits, e.g. time and/or temperature.
624
- 625 62. Activities in handling vials containing live biological agents, must be performed
626 in such a way to prevent the contamination of other products or egress of the live
627 agents into the work environment or the external environment. The viability of

628 such organisms and their biological classification should take into consideration
629 as part of the management of such risks.

630
631 63. Care should be taken in the preparation, printing, storage and application of
632 labels, including any specific text for patient-specific product of the contents on
633 the immediate and outer packaging.

634
635 In the case of autologous products, the unique patient identifier and the statement
636 “for autologous use only” should be indicated on the outer packaging or, where
637 there is no outer packaging, on the immediate packaging.

638
639 64. The compatibility of labels with ultra-low storage temperatures, where such
640 temperatures are used, should be verified.

641
642 65. Where donor (human or animal health) information becomes available after
643 procurement, which affects product quality, it should be taken into account in
644 recall procedures.

645
646

647 **Quality Control**

648

649 66. In-process controls have a greater importance in ensuring the consistency of the
650 quality of biological active substance and medicinal products than for
651 conventional products. In-process control testing should be performed at
652 appropriate stages of production to control those conditions that are important for
653 the quality of the finished product.

654

655 67. Where intermediates can be stored for extended periods of time (days, weeks or
656 longer), consideration should be given to the inclusion of finished product batches
657 made from materials held for their maximum in-process periods in the on-going
658 stability programme.

659

660 68. Certain types of cells (e.g. autologous cells) may be available in limited quantities
661 and, where allowed in the MA, a modified testing and sample retention strategy
662 may be developed and documented.

663

664 69. Forcellular products, sterility tests should be conducted on antibiotic-free cultures
665 of cells or cell banks to provide evidence for absence of bacterial and fungal
666 contamination and to be able to detection fastidious organisms where
667 appropriate.

668

669 70. For biological medicinal products with a short shelf life, which for the purposes of
670 the annex is taken to mean a period that does not permit release when sterility
671 testing results are provided after 14 days or less, and which need batch
672 certification before completion of all end product quality control tests (e.g. sterility
673 tests) a suitable control strategy must be in place. Such controls need to be built
674 on enhanced understanding of product and process performance and take into
675 account the controls and attributes of starting and raw materials. The exact and
676 detailed description of the entire release procedure, including the responsibilities
677 of the different personnel involved in assessment of production and analytical
678 data is essential. A continuous assessment of the effectiveness of the quality

679 assurance system must be in place including records kept in a manner which
680 permit trend evaluation.

681
682 Where end product tests are not available due to their short shelf life, alternative
683 methods of obtaining equivalent data to permit batch certification should be
684 considered (e.g. rapid microbiological methods). The procedure for batch
685 certification and release may be carried out in two or more stages:

686
687 (a) Assessment by designated person(s) of batch processing records, results
688 from environmental monitoring (where available) which should cover
689 production conditions, all deviations from normal procedures and the
690 available analytical results for review in preparation for the initial
691 certification by the Responsible Person.

692 (b) Assessment of the final analytical tests and other information available for
693 final certification by the Authorised Person. A procedure should be in place
694 to describe the measures to be taken (including liaison with clinical staff)
695 where out of specification test results are obtained. Such events should be
696 fully investigated and the relevant corrective and preventive actions taken
697 to prevent recurrence documented.

698
699

700 **PART B. SPECIFIC GUIDANCE ON SELECTED PRODUCT** 701 **TYPES**

702

703 **B1. ANIMAL SOURCED PRODUCTS²⁵**

704

705 This guidance applies to animal materials which includes materials from
706 establishments such as abattoirs. Since the supply chains can be extensive and
707 complex, controls based on QRM principles need to be applied, see also
708 requirements of appropriate pharmacopoeial monographs, including the need for
709 specific tests at defined stages. Documentation to demonstrate the supply chain
710 traceability²⁶ and clear roles of participants in the supply chain, typically including
711 a sufficiently detailed and current process map, should be in place.

712

713 1. Monitoring programmes should be in place for animal disease that are of concern
714 to human health. Organisations should take into account reports from trustworthy
715 sources on national disease prevalence when compiling their assessment of risk
716 and mitigation factors. Such organisations include the World Organisation for
717 Animal Health (OIE, Office International des Epizooties²⁷). This should be
718 supplemented by information on health monitoring and control programme(s) at
719 national and local levels, the latter to include the sources (e.g. farm or feedlot)
720 from which the animals are drawn and the control measures in place during
721 transport to the abattoirs.

722

723 2. Where abattoirs are used to source animal tissues, they should be shown to
724 operate to stringent standards. Account should be taken of reports from national

²⁵ In the EEA, see also PhEur monograph requirements, 0333

²⁶ See PIC/S GMP Chapter 5.

²⁷ http://www.oie.int/eng/en_index.htm

- 725 regulatory organisations²⁸ which verify compliance with the requirements of food,
726 safety, and quality veterinary and plant health legislation.
727
- 728 3. Control measures for starting or raw materials at establishments such as
729 abattoirs should include appropriate elements of a Quality Management System
730 to assure a satisfactory level of operator training, materials traceability, control
731 and consistency. These measures may be drawn from sources outside PIC/S
732 GMP but should be shown to provide equivalent levels of control.
733
- 734 4. Control measures for starting or raw materials should be in place which prevent
735 interventions which may affect the quality of materials, or which at least provides
736 evidence of such activities, during their progression through the manufacturing
737 and supply chain. This includes the movement of material between sites of initial
738 collection, partial and final purification(s), storage sites, hubs, consolidators and
739 brokers. Details of such arrangements should be recorded within the traceability
740 system and any breaches recorded, investigated and actions taken.
741
- 742 5. Regular audits of the starting or raw material supplier should be undertaken which
743 verify compliance with controls for materials at the different stages of
744 manufacture. Issues must be investigated to a depth appropriate to their
745 significance, for which full documentation should be available. Systems should
746 also be in place to ensure that effective corrective and preventive actions are
747 taken.
748

749

750 **B2. ALLERGEN PRODUCTS**

751

752 Materials may be manufactured by extraction from natural sources or
753 manufactured by recombinant DNA technology.

754

- 755 1. Source materials should be described in sufficient detail to ensure consistency in
756 their supply, e.g. common and scientific name, origin, nature, contaminant limits,
757 method of collection. Those derived from animals should be from healthy
758 sources. Appropriate biosecurity controls should be in place for colonies (e.g.
759 mites, animals) used for the extraction of allergens. Allergen products should be
760 stored under defined conditions to minimise deterioration.
761
- 762 2. The production process steps including pre-treatment, extraction, filtration,
763 dialysis, concentration or freeze-drying steps should be described in detail and
764 validated.
765
- 766 3. The modification processes to manufacture modified allergen extracts (e.g.
767 allergoids, conjugates) should be described. Intermediates in the manufacturing
768 process should be identified and controlled.
769
- 770 4. Allergen extract mixtures should be prepared from individual extracts from single
771 source materials. Each individual extract should be considered as one active
772 substance.
773

774

²⁸ In the EEA, this is the Food and Veterinary Office http://ec.europa.eu/food/fvo/index_en.htm.

775 **B.3 ANIMAL IMMUNOSERA PRODUCTS**

776

777 1. Particular care should be exercised on the control of antigens of biological origin
778 to assure their quality, consistency and freedom from adventitious agents. The
779 preparation of materials used to immunise the source animals (e.g. antigens,
780 haptens carriers, adjuvants, stabilising agents), the storage of such material
781 immediately prior to immunisation should be in accordance with documented
782 procedures.

783

784 2. The immunisation, test bleed and harvest bleed schedules should conform to
785 those approved in the CTA or MA.

786

787 3. The manufacturing conditions for the preparation of antibody sub-fragments (e.g.
788 Fab or F(ab')₂) and any further modifications must be in accordance with
789 validated and approved parameters. Where such enzymes are made up of
790 several components, their consistency should be assured.

791

792

793 **B.4 VACCINES**

794

795 1. Where eggs are used, the health status of all source flocks used in the production
796 of eggs (whether specified pathogen free or healthy flocks) should be assured.

797

798 2. The integrity of containers used to store intermediate products and the hold times
799 must be validated.

800

801 3. Vessels containing inactivated products should not be opened or sampled in
802 areas containing live biological agents.

803

804 4. The sequence of addition of active ingredients, adjuvants and excipients during
805 the formulation of an intermediate or final product must be in compliance with
806 specifications.

807

808 5. Where organisms with a higher biological safety level (e.g. pandemic vaccine
809 strains) are to be used in manufacture or testing, appropriate containment
810 arrangements must be in place. The approval of such arrangements should be
811 obtained from the appropriate national authority(ies) and the approval documents
812 be available for verification.

813

814

815 **B.5 RECOMBINANT PRODUCTS**

816

817 1. Process condition during cell growth, protein expression and purification must be
818 maintained within validated parameters to assure a consistent product with a
819 defined range of impurities that is within the capability of the process to reduce
820 to acceptable levels. The type of cell used in production may require increased
821 measures to be taken to assure freedom from viruses. For production involving
822 multiple harvest, the period of continuous cultivation should be within specified
823 limits.

824

- 825 2. The purification processes to remove unwanted host cell proteins, nucleic acids,
826 carbohydrates, viruses and other impurities should be within defined validated
827 limits.
828
829

830 **B6. MONOCLONAL ANTIBODY PRODUCTS**

- 831
832 1. Monoclonal antibodies may be manufactured from murine hybridomas, human
833 hybridomas or by recombinant DNA technology. Control measures appropriate
834 to the different source cells (including feeder cells if used) and materials used to
835 establish the hybridoma / cell line should be in place to assure the safety and
836 quality of the product. It should be verified that these are within approved limits.
837 Freedom from viruses should be given particular emphasis. It should be noted
838 that data originating from products generated by the same manufacturing
839 technology platform may be acceptable to demonstrate suitability.
840
841 2. Criteria to be monitored at the end of a production cycle and for early termination
842 of production cycles should be verified that these are within approved limits.
843
844 3. The manufacturing conditions for the preparation of antibody sub-fragment (e.g.
845 Fab, F(ab')₂, scFv) and any further modifications (e.g. radio labelling, conjugation,
846 chemical linking) must be in accordance with validated parameters.
847

848

849 **B7. TRANSGENIC ANIMAL PRODUCTS**

850

851 Consistency of starting material from a transgenic source is likely to be more
852 problematic than is normally the case for non-transgenic biotechnology sources.
853 Consequently, there is an increased requirement to demonstrate batch-to-batch
854 consistency of product in all respects.
855

- 856 1. A range of species may be used to produce biological medicinal products, which
857 may be expressed into body fluids (e.g. milk) for collection and purification.
858 Animals should be clearly and uniquely identified and backup arrangements
859 should be put in place in the event of loss of the primary marker.
860
861 2. The arrangements for housing and care of the animals should be defined such
862 that they minimise the exposure of the animals to pathogenic and zoonotic
863 agents. Appropriate measures to protect the external environment should be
864 established. A health-monitoring programme should be established and all
865 results documented, any incident should be investigated and its impact on the
866 continuation of the animal and on previous batches of product should be
867 determined. Care should be taken to ensure that any therapeutic products used
868 to treat the animals do not contaminate the product.
869
870 3. The genealogy of the founder animals through to production animals must be
871 documented. Since a transgenic line will be derived from a single genetic founder
872 animal, materials from different transgenic lines should not be mixed.
873
874 4. The conditions under which the product is harvested should be in accordance
875 with CTA or MA conditions. The harvest schedule and conditions under which

876 animals may be removed from production should be performed according to
877 approved procedures and acceptance limits.

878
879

880 **B8. TRANSGENIC PLANT PRODUCTS**

881
882
883
884
885
886

Consistency of starting material from a transgenic source is likely to be more problematic than is normally the case for non-transgenic biotechnology sources. Consequently, there is an increased requirement to demonstrate batch-to-batch consistency of product in all respects.

887 1. Additional measures, over and above those given in Part A, may be required to
888 prevent contamination of master and working transgenic banks by extraneous
889 plant materials and relevant adventitious agents. The stability of the gene within
890 defined generation numbers should be monitored.

891
892 2. Plants should be clearly and uniquely identified, the presence of key plant
893 features, including health status, across the crop should be verified at defined
894 intervals through the cultivation period to assure consistency of yield between
895 crops.

896
897 3. Security arrangements for the protection of crops should be defined, wherever
898 possible, such that they minimise the exposure to contamination by
899 microbiological agents and cross-contamination with non-related plants.
900 Measures should be in place to prevent materials such as pesticides and
901 fertilisers from contaminating the product. A monitoring programme should be
902 established and all results documented, any incident should be investigated and
903 its impact on the continuation of the crop in the production programme should be
904 determined.

905
906 4. Conditions under which plants may be removed from production should be
907 defined. Acceptance limits should be set for materials (e.g. host proteins) that
908 may interfere with the purification process. It should be verified that the results
909 are within approved limits.

910
911 5. Environmental conditions (temperature, rain), which may affect the quality
912 attributes and yield of the recombinant protein from time of planting, through
913 cultivation to harvest and interim storage of harvested materials should be
914 documented. The principles in documents such as 'Guideline on Good
915 Agricultural and Collection Practice for Starting Materials of Herbal Origin'²⁹
916 should be taken into account when drawing up such criteria.

917
918

919 **GLOSSARY**

920
921
922

See Annex 2A

²⁹ EMA, WHO or equivalent